

MATHEMATICAL NOTES.

By A. RÉNYI in Budapest.

II. On the sequence of generalized partial sums of a series.

Introduction.

Let $a_0, a_1, \dots, a_n, \dots$ denote an arbitrary sequence of real numbers. We define the corresponding sequence $A_0, A_1, \dots, A_n, \dots$ as follows: we put

$$A_0 = 0$$

and if the representation of the integer $n \geq 1$ in the dyadic system is

$$(1) \quad n = 2^{k_1} + 2^{k_2} + \dots + 2^{k_l}$$

(where $k_1 > k_2 > \dots > k_l \geq 0$ are integers) we put

$$(2) \quad A_n = a_{k_1} + a_{k_2} + \dots + a_{k_l}.$$

We shall call the sequence $\{A_n\}$ *the sequence of generalized partial sums of the sequence $\{a_n\}$ (or of the series $\sum_{n=0}^{\infty} a_n$)*.

Clearly the sequence $\{A_n\}$ consists of all possible finite sums of elements of the sequence $\{a_n\}$, each such sum occurring exactly once in the sequence $\{A_n\}$; the mentioned sums are ordered according to the lexicographic order. Evidently the ordinary partial sums of any rearrangement of the series $\sum_{n=0}^{\infty} a_n$ are all contained in the sequence $\{A_n\}$. Clearly

if the series $\sum_{n=0}^{\infty} a'_n$ is a rearrangement of the series $\sum_{n=0}^{\infty} a_n$, then the sequence $\{A'_n\}$ corresponding to the sequence $\{a'_n\}$ in the same way as $\{A_n\}$ corresponds to $\{a_n\}$, is a rearrangement of $\{A_n\}$. This is worth mentioning because by rearranging a series the sequence of ordinary partial sums is in general completely changed.

In the present paper we shall investigate how the properties of the sequence $\{A_n\}$ depend on the properties of the sequence $\{a_n\}$.

§ 1. On the limit of the arithmetic means of the powers of generalized partial sums.

First we prove the following

Theorem 1. *The limit*

$$\alpha_1 = \lim_{n \rightarrow \infty} \frac{\sum_{k=0}^{n-1} A_k}{n}$$

exists if and only if the series $\sum_{j=0}^{\infty} a_j$ converges; in this case we have

$$\alpha_1 = \frac{1}{2} \sum_{j=0}^{\infty} a_j.$$

PROOF OF THEOREM 1. Let us put

$$(3) \quad \sigma_n = \frac{1}{n} \sum_{k=0}^{n-1} A_k.$$

As clearly

$$(4) \quad A_k + A_{2^{s-1}-k} = \sum_{j=0}^{s-1} a_j \quad 0 \leq k \leq 2^s - 1$$

we have

$$(5) \quad \sigma_{2^s} = \frac{1}{2} \sum_{j=0}^{s-1} a_j.$$

Thus if $\alpha_1 = \lim_{n \rightarrow \infty} \sigma_n$ exists, we have also $\lim_{s \rightarrow \infty} \sigma_{2^s} = \alpha_1$ and thus $\sum_{j=0}^{\infty} a_j$ is convergent and has the sum $2\alpha_1$. This proves that the convergence of $\sum_{j=0}^{\infty} a_j$ is necessary for the existence of the limit α_1 . Now let us assume that $\sum_{j=0}^{\infty} a_j$ is convergent, and let us put

$$(6) \quad \sum_{j=0}^{\infty} a_j = A.$$

It follows by (5) that

$$(7) \quad \lim_{s \rightarrow \infty} \sigma_{2^s} = \frac{A}{2}.$$

It is easy to verify the following assertion: If $n = 2^{k_1} + 2^{k_2} + \dots + 2^{k_l}$ where

$k_1 > k_2 > \dots > k_l \geq 0$, we have

$$(8) \quad \sigma_n = \frac{\sum_{\nu=1}^l 2^{k_\nu} \sigma_{2^{k_\nu}}}{\sum_{\nu=1}^l 2^{k_\nu}} + \frac{\sum_{\nu=2}^l 2^{k_\nu} (a_{k_1} + a_{k_2} + \dots + a_{k_{\nu-1}})}{\sum_{\nu=1}^l 2^{k_\nu}}$$

Now we need the following elementary

Lemma 1. *If $c_0, c_1, \dots, c_k, \dots$ is an arbitrary sequence of numbers, with $\lim_{k \rightarrow \infty} c_k = c$, and we put*

$$(9) \quad \gamma_n = \frac{\sum_{\nu=1}^l 2^{k_\nu} c_{k_\nu}}{\sum_{\nu=1}^l 2^{k_\nu}}$$

for $n = 2^{k_1} + 2^{k_2} + \dots + 2^{k_l}$, then $\lim_{n \rightarrow \infty} \gamma_n = c$.

PROOF OF LEMMA 1. It is easy to see, that the linear summation method by which we obtain $\{\gamma_n\}$ from $\{c_k\}$ is a regular TOEPLITZ method; this proves Lemma 1. A direct proof is as follows:

Let us choose an arbitrary $\varepsilon > 0$; then there can be found an integer $K_0 = K_0(\varepsilon)$ such that $|c_k - c| < \varepsilon$ for $k \geq K_0$; we have further $|c_k| \leq C$ for $k = 0, 1, \dots$. Now we have clearly

$$|\gamma_n - c| \leq \varepsilon + \frac{C \sum_{k=0}^{K_0-1} 2^k}{n} \leq 2\varepsilon$$

for $n \geq \frac{C2^{K_0}}{\varepsilon}$. Thus $\lim_{n \rightarrow \infty} \gamma_n = c$.

It follows by Lemma 1 and (7) that the first term on the right of (8) tends to $\frac{A}{2}$. As regards the second term, we have clearly

$$(10) \quad \left| \frac{\sum_{\nu=2}^l 2^{k_\nu} (a_{k_1} + \dots + a_{k_{\nu-1}})}{\sum_{\nu=1}^l 2^{k_\nu}} \right| \leq \frac{\sum_{\nu=1}^l |a_{k_\nu}| \cdot 2^{k_\nu}}{\sum_{\nu=1}^l 2^{k_\nu}}$$

As $|a_n| \rightarrow 0$, it follows by Lemma 1 and (10) that the second term on the right of (8) tends to 0. Thus we have proved $\lim_{n \rightarrow \infty} \sigma_n = \frac{A}{2}$ and so the proof of Theorem 1 is completed.

Now we proceed to prove the following more general

Theorem 2. *The limits*

$$\alpha_r = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} A_k^r \quad (r = 1, 2, \dots)$$

all exist if and only if the series $\sum_{j=0}^{\infty} a_j$ and $\sum_{j=0}^{\infty} a_j^2$ are both convergent. The values of the limits α_r can be expressed as follows: let t denote a real number, $0 \leq t < 1$, and let us consider the dyadic expansion¹⁾

$$t = \sum_{n=1}^{\infty} \frac{\varepsilon_n(t)}{2^n}$$

of t , where $\varepsilon_n(t)$ is equal to 0 or 1. Let us consider the function

$$(11) \quad A(t) = \sum_{n=0}^{\infty} a_n \varepsilon_{n+1}(t).$$

If $\sum_{j=0}^{\infty} a_j$ and $\sum_{j=0}^{\infty} a_j^2$ are convergent, then the series on the right of (11) is convergent²⁾ for almost every value of t and the function $A(t)$ belongs to any class L^p ($p \geq 1$) in $(0, 1)$. The limits α_r are simply the moments of $A(t)$, i. e.

$$(12) \quad \alpha_r = \int_0^1 (A(t))^r dt \quad (r = 1, 2, \dots).$$

Before proving Theorem 2 we make some remarks.

Remark 1. It is clear from Theorem 1 that for the existence of α_1 the convergence of $\sum_{n=0}^{\infty} a_n^2$ is not necessary, but this condition is necessary already for the existence of α_2 . It is also clear that under the conditions of Theorem 2 we

have $\int_0^1 A(t) dt = \frac{A}{2}$, because $\int_0^1 \varepsilon_k(t) dt = \frac{1}{2}$ ($k = 1, 2, \dots$).

Remark 2. It follows from (12) for $r = 2$ that

$$\alpha_2 = \frac{\left(\sum_{j=0}^{\infty} a_j\right)^2 + \sum_{j=0}^{\infty} a_j^2}{4}.$$

¹⁾ If t is a dyadic rational number, $t = \frac{r}{2^s}$, we choose the finite expansion, in which $\varepsilon_n(t) = 0$ for $n > s$.

²⁾ See H. POLLARD, Subseries of a convergent series, *Bull. Amer. Math. Soc.* **49** (1943), 730–731.

This can be expressed also in the following form:

$$(13) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} (A_k - \alpha_1)^2 = \frac{1}{4} \sum_{j=0}^{\infty} a_j^2.$$

Remark 3. The functions $\varepsilon_k(t)$ are connected with the well known RADEMACHER functions $R_k(t) = \text{sg} \sin 2^k \pi t$ as follows: $\varepsilon_k(t) = \frac{1}{2}(1 + R_k(t))$. Thus (11) can be written also in the following equivalent form:

Let us put

$$(14) \quad D(t) = \frac{1}{2} \sum_{n=0}^{\infty} a_n R_{n+1}(t)$$

Then we have

$$(15) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} (A_k - \alpha_1)^r = \int_0^1 (D(t))^r dt \quad (r = 1, 2, \dots)$$

Clearly (13) is the special case $r = 2$ of (15). The right hand side of (15) is evidently equal to 0 for odd values of r .

Remark 4. Note that if $\sum_{j=0}^{\infty} a_j$ and $\sum_{j=0}^{\infty} a_j^2$ are convergent but $\sum_{j=0}^{\infty} |a_j|$ diverges, the sequence A_n is unbounded, and thus not only the existence of the limits α_r , but even the boundedness of the mean values $\frac{1}{n} \sum_{k=0}^{n-1} A_k^r$ is not trivial.

PROOF OF THEOREM 2. We start from the formula

$$(16) \quad \frac{1}{2^{\nu}} \sum_{k=0}^{2^{\nu}-1} A_k^r = \int_0^1 (S_{\nu}(t))^r dt$$

where

$$(17) \quad S_{\nu}(t) = \sum_{k=0}^{\nu-1} a_k \varepsilon_{k+1}(t).$$

To prove (16) it suffices to point out that the values of the function $S_{\nu}(t)$ are the numbers $A_0, A_1, \dots, A_{2^{\nu}-1}$, and each of these values is taken on by the function $S_{\nu}(t)$ on a subinterval of length $\frac{1}{2^{\nu}}$.

We prove first the sufficiency part of Theorem 2.

Let us suppose that the series $\sum_{j=0}^{\infty} a_j$ and $\sum_{j=0}^{\infty} a_j^2$ are convergent and put $\sum_{j=0}^{\infty} a_j = A$ and $\sum_{j=0}^{\infty} a_j^2 = B^2$. By a well known theorem on RADEMACHER'S

series³⁾ the convergence of the series $\sum_{k=0}^{\infty} b_k^2$ implies that the series $\sum_{k=0}^{\infty} b_k R_k(t)$ converges almost everywhere to a function which belongs to every class L^p ($p \geq 1$). It is also known that for any integer $m \geq 1$ we have

$$(18) \quad \int_0^1 \left| \sum_{k=0}^N b_k R_k(t) \right|^{2m} dt \leq C_m \left(\sum_{k=0}^N b_k^2 \right)^m$$

where C_m is a positive constant, not depending on b_0, b_1, \dots . We may take e. g. $C_m = m^m$. It follows that we have for almost every t

$$\lim_{\nu \rightarrow \infty} S_\nu(t) = A(t)$$

and $A(t)$ belongs to every class L^p ($p \geq 1$). We have further for $r \geq 2$

$$(19) \quad \left| \int_0^1 (S_\nu(t))^r dt - \int_0^1 (A(t))^r dt \right| \leq \int_0^1 |S_\nu(t)^r - A(t)^r| dt.$$

As for any pair of real numbers x and h we have for $r = 1, 2, \dots$

$$|(x+h)^r - x^r| \leq rh(|x|^{r-1} + |x+h|^{r-1})$$

it follows from (19) that

$$(20) \quad \left| \int_0^1 (S_\nu(t))^r dt - \int_0^1 (A(t))^r dt \right| \leq r \int_0^1 |S_\nu(t) - A(t)| (|S_\nu(t)|^{r-1} + |A(t)|^{r-1}) dt.$$

Applying the inequality of SCHWARZ we obtain

$$(21) \quad \left| \int_0^1 (S_\nu(t))^r dt - \int_0^1 (A(t))^r dt \right| \leq 2r \left[\left(\int_0^1 (S_\nu(t) - A(t))^2 dt \right) (r-1)^{r-1} \left(\frac{A^2 + B^2}{4} \right)^{r-1} \right]^{\frac{1}{2}}$$

because by (18)

$$\int_0^1 |S_\nu(t)|^{2r-2} dt \leq (r-1)^{r-1} \left(\frac{A^2 + B^2}{4} \right)^{r-1}$$

and

$$\int_0^1 (A(t))^{2r-2} dt \leq (r-1)^{r-1} \left(\frac{A^2 + B^2}{4} \right)^{r-1}.$$

Now as

$$\int_0^1 (S_\nu(t) - A(t))^2 dt = \frac{\sum_{k=\nu}^{\infty} a_k^2 + \left(\sum_{k=\nu}^{\infty} a_k \right)^2}{4},$$

³⁾ See A. ZYGMUND, Trigonometrical Series, Monografie Matematyczne, Warszawa—Lwów 1935, pp. 123—124.

it follows from (21) that

$$\lim_{r \rightarrow \infty} \int_0^1 (S_r(t))^r dt = \int_0^1 (A(t))^r dt.$$

Thus putting

$$(22) \quad \sigma_n^{(r)} = \frac{1}{n} \sum_{k=0}^{n-1} A_k^r$$

we have proved

$$(23) \quad \lim_{r \rightarrow \infty} \sigma_{2^r}^{(r)} = \int_0^1 (A(t))^r dt = \alpha_r.$$

To deduce $\lim_{n \rightarrow \infty} \sigma_n^{(r)} = \alpha_r$ from (23), we shall need the following identity:

If $n = 2^{k_1} + 2^{k_2} + \dots + 2^{k_l}$ with $k_1 > k_2 > \dots > k_l \geq 0$ we have

$$(24) \quad \sum_{k=0}^{n-1} A_k^r = \sum_{j=1}^l 2^{k_j} \sigma_{2^{k_j}}^{(r)} + \sum_{j=2}^l \left[\sum_{i=0}^{2^{k_j}-1} \left(A_i + \sum_{h=1}^{j-1} a_{k_h} \right)^r - A_i^r \right]$$

Thus it follows

$$(25) \quad \sigma_n^{(r)} = u_n^{(r)} + \sum_{\varrho=0}^{r-1} \binom{r}{\varrho} v_n^{(r, \varrho)},$$

where

$$(26) \quad u_n^{(r)} = \frac{\sum_{j=1}^l 2^{k_j} \sigma_{2^{k_j}}^{(r)}}{\sum_{j=1}^l 2^{k_j}}$$

and

$$(27) \quad v_n^{(r, \varrho)} = \frac{\sum_{j=2}^l 2^{k_j} \sigma_{2^{k_j}}^{(\varrho)} (a_{k_1} + \dots + a_{k_{j-1}})^{r-\varrho}}{\sum_{j=1}^l 2^{k_j}}$$

It follows by (23) and Lemma 1 that

$$(28) \quad \lim_{n \rightarrow \infty} u_n^{(r)} = \alpha_r.$$

As regards $v_n^{(r, \varrho)}$ we shall prove that

$$(29) \quad \lim_{n \rightarrow \infty} v_n^{(r, \varrho)} = 0 \text{ for } \varrho = 0, 1, \dots, r-1; r = 1, 2, \dots$$

As every convergent sequence is bounded, there can be found positive constants K_ϱ ($\varrho = 1, 2, \dots$) such that

$$(30) \quad |\sigma_{2^v}^{(\varrho)}| \leq K_\varrho.$$

It will be useful to put $\sigma_{2^v}^{(0)} \equiv 1$ and thus $K_0 = 1$. It follows

$$(31) \quad |v_n^{(r, \varrho)}| \leq K_\varrho \frac{\sum_{j=2}^l 2^{kj} (|a_{k_1}| + \dots + |a_{k_{j-1}}|)^{r-\varrho}}{\sum_{j=1}^l 2^{kj}}.$$

Applying the inequality of CAUCHY, we obtain

$$(32) \quad |v_n^{(r, \varrho)}| \leq \frac{K_\varrho}{n} \sum_{j=2}^l 2^{kj} \left(\sum_{i=1}^{j-1} a_{k_i}^2 \right)^{\frac{r-\varrho}{2}} j^{\frac{r-\varrho}{2}}$$

Now to an arbitrary $\varepsilon > 0$ there can be found an integer $k_0 = k_0(\varepsilon)$ such that $\sum_{k=k_0}^{\infty} a_k^2 < \varepsilon^2$. We may further suppose $B^2 = \sum_{k=0}^{\infty} a_k^2 < 1$, because if this were not so, we could consider the sequence $\{\mathcal{G}a_n\}$ with a suitable \mathcal{G} ($0 < \mathcal{G} < 1$) instead of the sequence $\{a_n\}$. Thus it follows, taking into account that $k_j \leq k_1 - j$, that

$$(33) \quad |v_n^{(r, \varrho)}| \leq K_\varrho \left(\varepsilon \sum_{j=1}^{\infty} \frac{j^{\frac{r-\varrho}{2}}}{2^j} + \frac{2^{k_0} l^{\frac{r-\varrho}{2}}}{n} \right),$$

the series $\sum_{j=1}^{\infty} \frac{j^\alpha}{2^j}$ being convergent for any $\alpha > 0$.

As $l \leq k_1 + 1 \leq \frac{\log 2n}{\log 2}$ we obtain

$$(34) \quad |v_n^{(r, \varrho)}| \leq C_r \left(\varepsilon + \frac{2^{k_0} \left(\frac{\log 2n}{\log 2} \right)^{\frac{r-\varrho}{2}}}{n} \right) \quad (\varrho = 0, 1, \dots, r-1)$$

where C_r is a positive constant, depending only on r . Thus it follows that

$$(35) \quad \lim_{n \rightarrow \infty} v_n^{(r, \varrho)} = 0 \quad (0 \leq \varrho \leq r-1; r = 1, 2, \dots).$$

This completes the proof of $\lim_{n \rightarrow \infty} \sigma_n^{(r)} = \alpha_r$, and thus the proof of the sufficiency of the conditions of Theorem 2.

To prove the necessity it suffices to mention, that according to Theorem 1 the convergence of $\sum_{k=0}^{\infty} a_k$ is necessary already for the existence of α_1 , and

according to (16)

$$\sigma_{2^v}^{(2)} = \frac{\left(\sum_{k=0}^{v-1} a_k\right)^2 + \sum_{k=0}^{v-1} a_k^2}{4}.$$

Thus $\lim_{v \rightarrow \infty} \sigma_{2^v}^{(2)} = \alpha_2$ implies the convergence of $\sum_{k=0}^{\infty} a_k^2$.

It follows evidently from Theorem 2 that we have under the conditions of Theorem 2

$$(36) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} f(A_k) = \int_0^1 f(A(t)) dt$$

if $f(x)$ is any polynomial. If $\sum_{n=0}^{\infty} |a_n|$ is also convergent, i. e. if the sequence A_k is bounded, it follows by a well known argument that (36) is valid for any continuous function. If the absolute convergence of $\sum_{n=0}^{\infty} a_n$ is not supposed, we can prove (36) only for continuous functions $f(x)$ satisfying some restrictions concerning the order of magnitude of $f(x)$ for $x \rightarrow \infty$. We shall not go into details here, and mention only that without any restriction on $f(x)$ the integral on the right-hand side of (36) does not exist in general.⁴⁾

For some special continuous functions $f(x)$ the validity of (36) can be deduced from Theorem 2. For instance if $f(x) = |x - c|$ where c is an arbitrary real number, then (36) is valid. This implies, that

$$(37) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} |A_k - a_1| = \int_0^1 |D(t)| dt$$

and as the right-hand side (37) is positive unless $a_0 = a_1 = \dots = 0$, it follows from (37) that the sequence A_n is not strongly summable except if all a_n vanish.

⁴⁾ A sufficient condition for the existence of the integral on the right of (36) is that $f(x)$ should satisfy the inequality $f(x) \leq C e^{\delta x^2}$ where $\delta \leq \frac{4}{e} (A^2 + B^2)^{-1}$ (see ZYGMUND, loc. cit.).

§ 2. The asymptotic distribution of generalized partial sums.

Now we shall consider the asymptotic distribution of the sequence A_n . We shall prove the following

Theorem 3. Let $a_0, a_1, \dots, a_n, \dots$ denote a real sequence and put $A_0 = 0$ and $A_n = a_{k_1} + a_{k_2} + \dots + a_{k_l}$ if $n = 2^{k_1} + 2^{k_2} + \dots + 2^{k_l}$ where $k_1 > k_2 > \dots > k_l \geq 0$ are integers. Let $N_n(x)$ denote the number of those among the numbers A_0, A_1, \dots, A_{n-1} which are $< x$, i. e. put

$$(38) \quad N_n(x) = \sum_{\substack{A_k < x \\ k < n}} 1 \quad (n = 1, 2, \dots).$$

Let us suppose that the series $\sum_{j=0}^{\infty} a_j$ and $\sum_{j=0}^{\infty} a_j^2$ are convergent, and let us put again $A(t) = \sum_{n=0}^{\infty} a_n \varepsilon_{n+1}(t)$ where $\varepsilon_n(t)$ is the n -th dyadic digit of t , i. e. $\varepsilon_n(t)$ is 0 or 1 and $t = \sum_{n=1}^{\infty} \frac{\varepsilon_n(t)}{2^n}$. Let $F(x)$ denote the measure of the set of those points t of the interval $(0, 1)$ for which $A(t) < x$. By other words $F(x)$ is the distribution function of $A(t)$. ($F(x)$ is clearly nondecreasing, continuous to the right, $\lim_{x \rightarrow -\infty} F(x) = 0$ and $\lim_{x \rightarrow +\infty} F(x) = 1$.) Then we have

$$(39) \quad \lim_{n \rightarrow \infty} \frac{N_n(x)}{n} = F(x)$$

in all continuity points x of $F(x)$.

PROOF OF THEOREM 3. First we prove that if x is a point of continuity of $F(x)$ then

$$(40) \quad \lim_{v \rightarrow \infty} \frac{N_{2^v}(x)}{2^v} = F(x).$$

This can be shown as follows: The function

$$S_v(t) = \sum_{k=0}^{v-1} a_k \varepsilon_{k+1}(t)$$

takes on the values $A_0, A_1, \dots, A_{2^v-1}$, each on a set of measure $\frac{1}{2^v}$. Thus denoting by $F_v(x)$ the measure of the set of those values t ($0 \leq t < 1$) for which $S_v(t) < x$ (i. e. $F_v(x)$ is the distribution function of $S_v(t)$) we have

$$(41) \quad \frac{N_{2^v}(x)}{2^v} = F_v(x).$$

Taking into account that if $\sum_{j=0}^{\infty} a_j$ and $\sum_{j=0}^{\infty} a_j^2$ are convergent, $S_\nu(t)$ tends almost everywhere to $A(t)$, and therefore it tends also in measure to $A(t)$, we have $\lim_{\nu \rightarrow \infty} F_\nu(x) = F(x)$ in all continuity points of $F(x)$. This proves (40).

To prove (39) we start from the formula

$$(42) \quad N_n(x) = 2^{k_1} F_{2^{k_1}}(x) + \sum_{j=2}^l 2^{k_j} F_{2^{k_j}}(x - a_{k_1} - \dots - a_{k_{j-1}})$$

if

$$n = 2^{k_1} + 2^{k_2} + \dots + 2^{k_l} \quad (k_1 > k_2 > \dots > k_l \geq 0).$$

Now we need the following elementary

Lemma 2. *If $c_n \geq 0$ and $c_n \rightarrow 0$, there can be found a monotonically increasing sequence l_n of integers, such that $\lim_{n \rightarrow \infty} l_n = +\infty$ and*

$$\lim_{n \rightarrow \infty} (c_{n-1} + \dots + c_{n-l_n}) = 0.$$

PROOF OF THE LEMMA. Let us put

$$d_n = \text{Max}_{k \geq n} c_k$$

and

$$l_n = \min \left([n/2], \frac{1}{\sqrt{d_{[n/2]}}} \right).$$

Then we have

$$\sum_{j=n-l_n}^{n-1} c_j \leq \sum_{j=n-l_n}^{n-1} d_j \leq l_n d_{n-l_n} \leq \frac{d_{[n/2]}}{\sqrt{d_{[n/2]}}} = \sqrt{d_{[n/2]}} \rightarrow 0,$$

which proves the assertion of our Lemma.

To complete the proof of Theorem 3 let us choose the sequence l_n in such a way, that $l_n \rightarrow \infty$ for $n \rightarrow \infty$ and $(|a_\nu| + |a_{\nu-1}| + \dots + |a_{\nu-l_\nu}|) \rightarrow 0$ for $\nu \rightarrow \infty$. This is possible by Lemma 2 because $\sum_{j=0}^{\infty} a_j$ being convergent we have $\lim_{n \rightarrow \infty} a_n = 0$. For any $\varepsilon > 0$ we can find an integer $k^* = k^*(\varepsilon)$ such that $|F_{2^k}(x) - F(x)| < \varepsilon$ if $k \geq k^*$. It follows that

$$(43) \quad \frac{N_n(x)}{n} \leq \frac{1}{n} \sum_{j=1}^l 2^{k_j} F_{2^{k_j}}(x) \leq F(x) + \varepsilon + \frac{2^{k^*}}{n}$$

which implies that

$$(44) \quad \limsup_{n \rightarrow \infty} \frac{N_n(x)}{n} \leq F(x).$$

Clearly we can find a value ν_0 such that if $\nu \geq \nu_0$ then $|a_\nu| + |a_{\nu-1}| + \dots + |a_{\nu-l_\nu}| < \varepsilon$. Thus if $n \geq 2^{\nu_0}$ we have $k_1 \geq \nu_0$ and

$$\sum_{k_i \geq k_1 - l_{k_1}} 2^{k_i} \geq \sum_{i=1}^l 2^{k_i} - \sum_{r=0}^{k_1 - l_{k_1} - 1} 2^r \geq n - 2^{k_1 - l_{k_1}}.$$

We have evidently

$$\frac{N_n(x)}{n} \geq \frac{1}{n} \sum_{k_i > k_1 - l_{k_1}} 2^{k_i} F_{2^{k_i}}(x - \varepsilon).$$

If we choose the integer k^{**} so that for $k \geq k^{**}$ we have $|F_{2^k}(x - \varepsilon) - F(x - \varepsilon)| < \varepsilon$, it follows

$$(45) \quad \frac{N_n(x)}{n} \geq (F(x - \varepsilon) - \varepsilon) \left(1 - \frac{1}{2^{l_{k_1}}}\right)$$

provided that $k_1 - l_{k_1} \geq k^{**}$. As for $n \rightarrow \infty$, $k_1 \rightarrow \infty$ and thus $l_{k_1} \rightarrow \infty$, further $k_1 - l_{k_1} \rightarrow \infty$, it follows from (45) that

$$(46) \quad \liminf_{n \rightarrow \infty} \frac{N_n(x)}{n} \geq F(x).$$

(44) and (46) together imply that

$$(47) \quad \lim_{n \rightarrow \infty} \frac{N_n(x)}{n} = F(x)$$

for any continuity point x of $F(x)$. Thus Theorem 3 is proved.

§ 3. Equivalence of the (C, 2)-summability of generalized partial sums with the convergence of a series.

It follows from Theorem 1 that if the arithmetic means of the sequence $\{A_k\}$ of generalized partial sums of a series $\sum_{n=0}^{\infty} a_n$ converge to a limit α_1 , the series itself is convergent and has the sum $2\alpha_1$. In this § we shall show that the same holds for the Cesàro means of order 2 too. Thus we prove the following

Theorem 4. *If the Cesàro means of order 2 of the sequence $\{A_n\}$ of generalized partial sums of the series $\sum_{n=0}^{\infty} a_n$ converge to a limit α_1 , then the series $\sum_{n=0}^{\infty} a_n$ is convergent and has the sum $2\alpha_1$.*

PROOF OF THEOREM 4. Let us put

$$(48) \quad \sigma_n^{(2)} = \frac{1}{\binom{n}{2}} \sum_{k=0}^{n-1} (n-k) A_k.$$

By supposition $\lim_{n \rightarrow \infty} \sigma_n^{(2)} = \alpha_1$, which implies that

$$(49) \quad \lim_{n \rightarrow \infty} \sigma_{2^v}^{(2)} = \alpha_1.$$

Now it is easy to show that, putting

$$(50) \quad S_k = a_0 + a_1 + \cdots + a_k,$$

we have

$$(51) \quad \sigma_{2^v}^{(2)} = \frac{(2^{v-1} - 1)S_{v-1} + \sum_{i=0}^{v-2} 2^i \cdot S_i}{2(2^v - 1)}.$$

It follows that

$$(52) \quad \frac{2(2^{v+1} - 1)\sigma_{2^{v+1}}^{(2)} - 2(2^v - 1)\sigma_{2^v}^{(2)}}{(2^v - 1)} = S_v + \frac{S_{v-1}}{2^v - 1}.$$

By (49), we have

$$(53) \quad \lim_{v \rightarrow \infty} \left(S_v + \frac{S_{v-1}}{2^v - 1} \right) = 2\alpha_1.$$

Now we shall prove that (53) implies

$$(54) \quad \lim_{v \rightarrow \infty} S_v = 2\alpha_1.$$

Clearly it suffices to show that S_v is bounded. But if S_v were unbounded, we could find a subsequence S_{v_j} such that $v_j \rightarrow \infty$, $|S_{v_j}| \rightarrow \infty$ and $|S_{v_j}| \cong \cong |S_{v_j-1}|$ which would imply $\limsup_{v \rightarrow \infty} \left| \left(S_v + \frac{S_{v-1}}{2^v - 1} \right) \right| = +\infty$ in contradiction to (53). Thus Theorem 4 is proved. Similar results hold for other methods of summation too. We hope to return to the question in another paper.

(Received December 12, 1956.)