

Frequently visited sets for random walks

Endre Csáki* Antónia Földes† Pál Révész* Jay Rosen‡ Zhan Shi

Abstract

We study the occupation measure of various sets for a symmetric transient random walk in Z^d with finite variances. Let $\mu_n^X(A)$ denote the occupation time of the set A up to time n . It is shown that $\sup_{x \in Z^d} \mu_n^X(x + A)/\log n$ tends to a finite limit as $n \rightarrow \infty$. The limit is expressed in terms of the largest eigenvalue of a matrix involving the Green's function of X restricted to the set A . Some examples are discussed and the connection to similar results for Brownian motion is given.

AMS 2000 Subject Classification: Primary 60G50; Secondary 60F15, 60J55

Keywords: random walk, occupation measure, strong theorems

1 Introduction

Let $X_n, n = 0, 1, \dots$ be a symmetric transient random walk in Z^d ($d \geq 3$). We will always assume that $X_n, n = 0, 1, \dots$ is not supported on any subgroup strictly smaller than Z^d . We denote by μ_n^X its *occupation measure*:

$$\mu_n^X(A) = \sum_{j=0}^n \mathbf{1}_A(X_j)$$

*Research supported by the Hungarian National Foundation for Scientific Research, Grant No. T 037886 and T 043037.

†Research supported by a PSC CUNY Grant, No. 65685-0034.

‡Research supported, in part, by grants from the NSF and from PSC-CUNY.

for all sets $A \subseteq Z^d$. Let $q_n(x) = \mathbf{P}(X_n = x)$. As usual, we let

$$G(x) = \sum_{k=0}^{\infty} q_k(x) \quad (1.1)$$

denote the Green's function for $\{X_n\}$. For any finite $A \subseteq Z^d$ let Λ_A denote the largest eigenvalue of the $|A| \times |A|$ matrix

$$G_A(x, y) = G(x - y), \quad x, y \in A. \quad (1.2)$$

Theorem 1.1 *If X has finite second moments then*

$$\lim_{n \rightarrow \infty} \sup_{x \in Z^d} \frac{\mu_n^X(x + A)}{\log n} = -1/\log(1 - 1/\Lambda_A) \quad \text{a.s.} \quad (1.3)$$

and

$$\lim_{n \rightarrow \infty} \sup_{0 \leq m \leq n} \frac{\mu_n^X(X_m + A)}{\log n} = -1/\log(1 - 1/\Lambda_A) \quad \text{a.s.} \quad (1.4)$$

For our first example, when $A = \{0\}$, $\Lambda_A = G(0) = 1/\gamma_d$, where γ_d is the probability of no-return to the origin, and in the case of the simple random walk we recover Theorem 13 of [3].

Here are some other examples. Set $t_y = \mathbf{P}(T_y < \infty)$, where $T_y := \inf\{s > 0 : X_s = y\}$. Let $S(0, 1) = \{e_1, \dots, e_d, -e_1, \dots, -e_d\}$, $B(0, 1) = \{0\} \cup S(0, 1)$, be the (Euclidean) sphere and ball in Z^d of radius 1 centered at the origin.

Theorem 1.2 *If X has finite second moments, then for any $0 \neq y \in Z^d$*

$$\lim_{n \rightarrow \infty} \sup_{x \in Z^d} \frac{\mu_n^X(x + \{0, y\})}{\log n} = -1/\log(1 - \gamma_d/(1 + t_y)) \quad \text{a.s.} \quad (1.5)$$

For the simple random walk

$$\lim_{n \rightarrow \infty} \sup_{x \in Z^d} \frac{\mu_n^X(x + S(0, 1))}{\log n} = -1/\log(1 - \gamma_d/2d(1 - \gamma_d)) \quad \text{a.s.} \quad (1.6)$$

and

$$\lim_{n \rightarrow \infty} \sup_{x \in Z^d} \frac{\mu_n^X(x + B(0, 1))}{\log n} = -1/\log\left(\frac{p + \sqrt{p^2 + 2/d}}{2}\right) \quad \text{a.s.} \quad (1.7)$$

where $p = 1 - 1/2d(1 - \gamma_d)$.

Corollary 1.3 *If X has finite second moments, then for any fixed $K > 0$*

$$\lim_{n \rightarrow \infty} \frac{\max_{x, y \in Z^d: |x-y| \leq K} (\mu_n^X(\{x, y\}))}{\log n} < -2/\log(1 - \gamma_d) \quad \text{a.s.}$$

Since the constant for one-point set in Theorem 1.1 is $-1/\log(1 - \gamma_d)$, this corollary expresses the fact that any two points with individual occupation measures up to time n , both close to the maximum, should be at a distance larger than any constant $K > 0$. In particular, a neighbor of a maximally visited point is not maximally visited.

Let W_t denote Brownian motion in \mathbb{R}^d , $d \geq 3$. We denote by ν_T^W its *occupation measure*:

$$\nu_T^W(A) = \int_0^T \mathbf{1}_A(W_t) dt$$

for all Borel sets $A \subseteq \mathbb{R}^d$. Let $K \subseteq \mathbb{R}^d$ be a fixed compact neighborhood of the origin which is the closure of its interior and set $K(x, r) = x + rK$.

As usual, we let

$$u^0(x) = \frac{c_d}{|x|^{d-2}} \quad (1.8)$$

denote the 0-potential density for $\{W_t\}$, where $c_d = 2^{-1}\pi^{-d/2}\Gamma(\frac{d}{2} - 1)$. Let Λ_K^0 denote the norm of

$$R_K f(x) = \int_K u^0(x - y) f(y) dy$$

considered as an operator from $L^2(K, dx)$ to itself. If $B(x, r)$ denotes the Euclidean ball in \mathbb{R}^d of radius r centered at x , it is known, [1], that $\Lambda_{B(0,1)}^0 = 2r_d^{-2}$ where r_d is the smallest positive root of the Bessel function $J_{d/2-2}$.

We mention that it can be shown, at least for K convex, that for any $S \in (0, \infty)$ and any $T \in (0, \infty]$,

$$\lim_{\epsilon \rightarrow 0} \sup_{|x| \leq S} \frac{\nu_T(K(x, \epsilon))}{\epsilon^2 |\log \epsilon|} = 2\Lambda_K^0 \quad \text{a.s.} \quad (1.9)$$

and

$$\lim_{\epsilon \rightarrow 0} \sup_{0 \leq t \leq T} \frac{\nu_T(K(W_t, \epsilon))}{\epsilon^2 |\log \epsilon|} = 2\Lambda_K^0 \quad \text{a.s.} \quad (1.10)$$

(These results are mentioned for motivation. They are not used in the rest of the paper).

For any $x \in R^d$ and $\epsilon > 0$, let $e_\epsilon(x) = x + [0, \epsilon]^d$, the cube of edglength ϵ with ‘lower’ corner at x . Set

$$\mathcal{L}_\epsilon(K) = \{x \in \epsilon Z^d \mid e_\epsilon(x) \subseteq K\}, \quad \text{and} \quad \mathcal{C}_\epsilon(K) = \bigcup_{x \in \mathcal{L}_\epsilon(K)} e_\epsilon(x) \quad (1.11)$$

and assume that

$$\lim_{\epsilon \rightarrow 0} \lambda^d(\mathcal{C}_\epsilon(K)) = \lambda^d(K) \quad (1.12)$$

where λ^d denotes Lebesgue measure.

Note that $\epsilon^{-1}\mathcal{L}_\epsilon(K) \subseteq Z^d$.

Theorem 1.4 *Assume that X_1 has $d-1$ moments and covariance matrix equal to the identity. Then*

$$\lim_{\epsilon \rightarrow 0} \epsilon^2 \Lambda_{\epsilon^{-1}\mathcal{L}_\epsilon(K)} = \Lambda_K^0 \quad (1.13)$$

and consequently

$$-\lim_{\epsilon \rightarrow 0} \epsilon^2 / \log(1 - 1/\Lambda_{\epsilon^{-1}\mathcal{L}_\epsilon(K)}) = \Lambda_K^0. \quad (1.14)$$

Section 2 states and proves the crucial Localization Lemma 2.2. Theorem 1.1 is proven in section 3, Theorem 1.2 and Corollary 1.3 are proven in section 4, and Theorem 1.4 is proven in section 5.

2 Localization for random walk occupation measures

We start by providing a convenient representation of the law of the total occupation measure $\mu_\infty^X(A)$. This representation is the counterpart of the Ciesielski-Taylor representation for the total occupation measure of spatial Brownian motion in [1, Theorem 1].

Let $(f, g)_A = \sum_{x \in A} f(x)g(x)$, and let $\delta_0(x)$ be the function on A defined by $\delta_0(x) = \delta(0, x)$, $x \in A$.

Lemma 2.1 *Let $\{X_n\}$ be a symmetric transient random walk in Z^d , and let A be a finite set in Z^d which contains the origin. Then,*

$$\mathbf{P}\left(\mu_\infty^X(A) > u\right) = \sum_j h_j \left(\frac{\lambda_j - 1}{\lambda_j}\right)^u \quad u = 0, 1, \dots, \quad (2.1)$$

where $\lambda_1 > \lambda_2 \geq \dots \geq \lambda_{|A|} \geq 1/2$ are the eigenvalues of the symmetric matrix G_A with the corresponding orthonormal eigenvectors $\phi_j(y)$, $h_j := (1, \phi_j)_A \phi_j(0)$.

Proof of Lemma 2.1: Let $\mathcal{J} = \mu_\infty^X(A)$ and set $\bar{G}(x-y) = \sum_{k=1}^\infty q_k(x-y) = G(x-y) - q_0(x-y)$.

Note that for any m ,

$$\begin{aligned} \mathbb{E}(\mathcal{J}^m) &= \mathbb{E}\left(\left\{\sum_{i=0}^\infty \mathbf{1}_A(X_i)\right\}^m\right) = \sum_{i_1, \dots, i_m=0}^\infty \mathbb{E}\left(\prod_{j=1}^m \mathbf{1}_A(X_{i_j})\right) \\ &= \sum_{k=1}^m \sum_{\substack{c_1, \dots, c_k \in [1, m] \\ c_1 + \dots + c_k = m}} \binom{m}{c_1, \dots, c_k} \sum_{A^k} \sum_{0 \leq n_1 < \dots < n_k < \infty} \prod_{j=1}^k q_{n_j - n_{j-1}}(x_j - x_{j-1}). \end{aligned} \quad (2.2)$$

Here, k is the number of distinct indices $n_1 < \dots < n_k$ among the indices i_1, \dots, i_m and c_l is the number of times that n_l appears, i.e. $c_l = \#\{j; 1 \leq j \leq m, i_j = n_l\}$. The factor $\binom{m}{c_1, \dots, c_k}$ is the number of ways to assign the value n_l to c_l of the indices i_1, \dots, i_m , for each $1 \leq l \leq k$.

Also, we have that

$$\begin{aligned} \sum_{A^k} \sum_{0 \leq n_1 < \dots < n_k < \infty} \prod_{j=1}^k q_{n_j - n_{j-1}}(x_j - x_{j-1}) &= \sum_{A^k} G(x_1) \prod_{j=2}^k \bar{G}(x_j - x_{j-1}) \\ &= (1, \bar{G}_A^{k-1} G_A \delta_0)_A. \end{aligned} \quad (2.3)$$

Hence (we justify the computations shortly)

$$\begin{aligned} \mathbb{E}(e^{\zeta \mathcal{J}}) &= 1 + \sum_{m=1}^\infty \frac{\zeta^m}{m!} \sum_{k=1}^m \sum_{\substack{c_1, \dots, c_k \in [1, m] \\ c_1 + \dots + c_k = m}} \binom{m}{c_1, \dots, c_k} (1, \bar{G}_A^{k-1} G_A \delta_0)_A \\ &= 1 + \sum_{k=1}^\infty \sum_{m=k}^\infty \sum_{\substack{c_1, \dots, c_k \in [1, m] \\ c_1 + \dots + c_k = m}} \prod_{j=1}^k \frac{\zeta^{c_j}}{c_j!} (1, \bar{G}_A^{k-1} G_A \delta_0)_A \end{aligned} \quad (2.4)$$

$$= 1 + \sum_{k=1}^{\infty} (e^{\zeta} - 1)^k (1, \bar{G}_A^{k-1} G_A \delta_0)_A$$

G_A is a symmetric matrix. Let $\psi(p)$ denote the characteristic function of X_1 . Then $\psi(p)$ is real and $|\psi(p)| \leq 1$. Thus $0 \leq 1 - \psi(p) \leq 2$, or equivalently $\frac{1}{1-\psi(p)} \geq \frac{1}{2}$. Hence, using the Fourier transform representation $G(x-y) = \int e^{i((x-y) \cdot p)} (1 - \psi(p))^{-1} dp$ we can see that $\sum_{x,y \in A} G_A(x,y) a_x a_y \geq \frac{1}{2} \sum_{x \in A} a_x^2$ for any $\{a_x \in \mathbb{R}^1; x \in A\}$. By the standard theory for symmetric matrices, G_A has all eigenvalues $\geq 1/2$, and the corresponding eigenvectors of G_A , denoted $\{\phi_j\}$ form an orthonormal basis of $L^2(A)$ (see [7, Theorems VI.15, VI.16]). Moreover, since the entries of G_A are strictly positive, by the Perron-Frobenius Theorem, see [8, Theorem XIII.43], the eigenspace corresponding to $\Lambda_A = \lambda_1$ is one dimensional, and we may and shall choose ϕ_1 such that $\phi_1(y) > 0$ for all $y \in A$.

Thus we can write (2.4) as

$$\begin{aligned} \mathbb{E} \left(e^{\zeta \mathcal{J}} \right) &= 1 + \sum_{k=1}^{\infty} (e^{\zeta} - 1)^k (1, \bar{G}_A^{k-1} G_A \delta_0)_A & (2.5) \\ &= 1 + \sum_{j=1}^{|A|} (1, \phi_j)_A (\phi_j, \delta_0)_A \sum_{k=1}^{\infty} (e^{\zeta} - 1)^k (\lambda_j - 1)^{k-1} \lambda_j \\ &= 1 + \sum_{j=1}^{|A|} h_j (e^{\zeta} - 1) \lambda_j \sum_{k=1}^{\infty} (e^{\zeta} - 1)^{k-1} (\lambda_j - 1)^{k-1} \end{aligned}$$

where $h_j = (1, \phi_j)_A (\phi_j, \delta_0)_A$. It is now easy to see that we can justify the derivation of (2.4) and (2.5) if

$$|(e^{\zeta} - 1)(\lambda_j - 1)| < 1, \quad \forall j. \quad (2.6)$$

In that case we can write (2.5) as

$$\mathbb{E} \left(e^{\zeta \mathcal{J}} \right) = 1 + \sum_j h_j \frac{(e^{\zeta} - 1) \lambda_j}{1 - (e^{\zeta} - 1)(\lambda_j - 1)}. \quad (2.7)$$

Since $\sum_{j=1}^{|A|} h_j = \sum_{j=1}^{|A|} (1, \phi_j)_A (\phi_j, \delta_0)_A = (1, \delta_0)_A = 1$ we have that

$$\mathbb{E} \left(e^{\zeta \mathcal{J}} \right) = \sum_j h_j \frac{e^{\zeta}}{1 - (e^{\zeta} - 1)(\lambda_j - 1)}. \quad (2.8)$$

Let $f_j = 1 - 1/\lambda_j = (\lambda_j - 1)/\lambda_j$. A straightforward calculation shows that

$$\frac{e^{\zeta}(1 - f_j)}{1 - e^{\zeta} f_j} = \frac{e^{\zeta}}{1 - (e^{\zeta} - 1)(\lambda_j - 1)} \quad (2.9)$$

so that

$$\mathbb{E} \left(e^{\zeta \mathcal{J}} \right) = \sum_j h_j \frac{e^{\zeta}(1 - f_j)}{1 - e^{\zeta} f_j}. \quad (2.10)$$

Note that since all $\lambda_j \geq 1/2$ we have $|f_j| \leq 1$. We can always choose ζ so that addition to (2.6) we also have

$$|e^{\zeta}| < 1. \quad (2.11)$$

Then we can write

$$\frac{e^{\zeta}(1 - f_j)}{1 - e^{\zeta} f_j} = e^{\zeta}(1 - f_j) \sum_{k=0}^{\infty} e^{k\zeta} f_j^k. \quad (2.12)$$

Hence

$$\sum_{k=1}^{\infty} e^{k\zeta} \mathbf{P}(\mathcal{J} = k) = \mathbb{E} \left(e^{\zeta \mathcal{J}} \right) = \sum_j h_j (1 - f_j) \sum_{k=1}^{\infty} e^{k\zeta} f_j^{k-1}. \quad (2.13)$$

We can choose $\zeta_0 < 0$ so that (2.6) and (2.11) hold. Furthermore, both sides of (2.13) are analytic functions of ζ in some neighborhood of $\zeta_0 + iR^1$ and agree for $\zeta_0 + iy$ when y is small. This is enough to allow us to conclude that

$$\mathbf{P}(\mathcal{J} = k) = \sum_j h_j (1 - f_j) f_j^{k-1}, \quad k = 1, 2, \dots \quad (2.14)$$

Hence

$$\mathbf{P}(\mathcal{J} > u) = \sum_j h_j f_j^u \quad u = 0, 1, \dots \quad (2.15)$$

This completes the proof of (2.1). \square

With the aid of (2.1) we next provide a localization result for the occupation measure of $\{X_n\}$.

Lemma 2.2 (The Localization Lemma) *Let $\{X_n\}$ be a symmetric transient random walk in Z^d with finite second moments, and let A be a finite set in Z^d . Set $\theta^* = \log(\Lambda_A/(\Lambda_A - 1))$. Then for some $c_1 < \infty$, $n \geq u^6$, and all $u > 0$ sufficiently large*

$$c_1^{-1} e^{-\theta^* u} \leq \mathbf{P} \left(\mu_n^X(A) \geq u \right) \leq \mathbf{P} \left(\mu_\infty^X(A) \geq u \right) \leq c_1 e^{-\theta^* u}. \quad (2.16)$$

Proof of Lemma 2.2: Let $\mathcal{J}_n := \mu_n^X(A)$. Assume first that A contains the origin. The dominant terms in (2.15) correspond to the f_j 's with largest absolute value. But since $\mathbf{P}(\mathcal{J} > u) \geq 0$ and monotone decreasing it is clear that these dominant terms must in addition be those which correspond to positive f_j 's. Thus the f_j 's with largest absolute value are positive, i.e. correspond to λ_j 's which are greater than 1. Recall that ϕ_1 is a strictly positive function on A , hence in (2.15) we have $h_1 > 0$. Since $(x-1)/x = 1 - 1/x$ is strictly monotone increasing on $(1, \infty)$ we conclude that the dominant term in (2.15) is precisely the single term corresponding to the largest eigenvalue $\lambda_1 = \Lambda_A$. Hence

$$\mathbf{P}(\mathcal{J} > u) \sim h_1 f_1^u = h_1 \left(\frac{\Lambda_A - 1}{\Lambda_A} \right)^u = h_1 e^{-u \log(\Lambda_A/(\Lambda_A - 1))} \quad (2.17)$$

implying that

$$\lim_{u \rightarrow \infty} \mathbf{P}(\mathcal{J}_\infty > u) e^{\theta^* u} = h_1 \in (0, \infty) \quad (2.18)$$

out of which the upper bound of (2.16) immediately follows.

Turning to prove the corresponding lower bound, let $\tau_z := \inf\{s : |X_s| > z\}$, and note that

$$\mathbf{P}(\tau_z > n) \leq c_1 \exp(-c_2 n z^{-2}). \quad (2.19)$$

Here is a simple proof:

$$\begin{aligned} \mathbf{P}(\tau_z > n) &= \mathbf{P}(|X_k| \leq z; 1 \leq k \leq n) \\ &\leq \mathbf{P}(|X_{lz^2}| \leq z; 1 \leq l \leq n z^{-2}) \\ &\leq \mathbf{P}(|X_{lz^2} - X_{(l-1)z^2}| \leq 2z; 1 \leq l \leq n z^{-2}) \\ &\leq \prod_{l=1}^{\lfloor n z^{-2} \rfloor} \mathbf{P}(|X_{lz^2} - X_{(l-1)z^2}| \leq 2z) \\ &= (\mathbf{P}(|X_{z^2}| \leq 2z))^{\lfloor n z^{-2} \rfloor} \leq e^{-c_2 \lfloor n z^{-2} \rfloor}. \end{aligned} \quad (2.20)$$

Hence

$$\mathbf{P}(\mathcal{J}_n > u) \geq \mathbf{P}(\mathcal{J}_{\tau_z} > u) - \mathbf{P}(\tau_z > n) \geq \mathbf{P}(\mathcal{J}_{\tau_z} > u) - c^{-1} \exp(-cnz^{-2}). \quad (2.21)$$

As usual we use the notation \mathbf{P}^a to denote probabilities of the random walk $a + X_n$, $n = 0, 1, \dots$. We now observe that

$$\sup_{a \in A} \mathbf{P}^a(\mu_\infty^X(A) > u) \leq c \mathbf{P}(\mu_\infty^X(A) > u) \quad (2.22)$$

for some $c < \infty$ and all u . To see this, note that for each $a \in A$ we can find some n_a with $h_a = \mathbf{P}(X_{n_a} = a) > 0$. Then using the Markov property,

$$\mathbf{P}(\mu_\infty^X(A) > u) \geq \mathbf{P}(\{\mu_\infty^X(A) > u\} \circ \theta_{n_a}, X_{n_a} = a) = h_a \mathbf{P}^a(\mu_\infty^X(A) > u). \quad (2.23)$$

Then (2.22) follows with $c = \sup_{a \in A} h_a^{-1} < \infty$.

Let \mathcal{J} and \mathcal{J}' denote two independent copies of \mathcal{J}_∞ and $T_A := \inf\{s > 0 : X_s \in A\}$. Noting (2.22), and using the strong Markov property, it is not hard to verify that

$$\mathbf{P}(\mathcal{J}_\infty > u) \leq \mathbf{P}(\mathcal{J}_{\tau_z} > u) + c \mathbf{P}(\mathcal{J} + \mathcal{J}' > u) \sup_{|v| > z} \mathbf{P}^v(T_A < \infty) \quad (2.24)$$

(c.f. [2, (3.6) and (3.7)] where this is obtained for the Brownian motion). It follows from Theorem 10.1 of [5] that

$$G(x) \leq \frac{c}{|x|}, \quad |x| \neq 0. \quad (2.25)$$

Using this together with the fact that $G(X_{n \wedge T_A})$ is a martingale shows that

$$G(v) = \mathbb{E}^v(G(X_{T_A}), T_A < \infty) \geq \inf_{a \in A} G(a) \mathbf{P}^v(T_A < \infty). \quad (2.26)$$

Therefore

$$\sup_{|v| > z} \mathbf{P}^v(T_A < \infty) \leq cz^{-1}. \quad (2.27)$$

By (2.18) it follows that for some constant C independent of u , which may change from line to line,

$$\mathbf{P}(\mathcal{J} + \mathcal{J}' > u) = \mathbf{P}(\mathcal{J} > u) + \sum_{y=0}^u \mathbf{P}(\mathcal{J}' > u - y) \mathbf{P}(\mathcal{J} = y)$$

$$\begin{aligned}
&\leq C \left[\exp(-u\theta^*) + \sum_{y=0}^u \exp(-(u-y)\theta^*) \mathbf{P}(\mathcal{J} = y) \right] \\
&\leq C \exp(-u\theta^*) + C \sum_{y=0}^u \exp(-u\theta^*) \\
&= C(1+u) \exp(-u\theta^*). \tag{2.28}
\end{aligned}$$

Hence, taking $z = u^2$ one gets from (2.21) and (2.24) that for some $c' > 0$, all $n \geq u^6$ and u sufficiently large

$$\mathbf{P}(\mathcal{J}_n > u) \geq c' e^{-\theta^* u} \tag{2.29}$$

as needed to complete the proof of the lemma when A contains the origin. In general we have

$$\mathbf{P}(\mu_\infty^X(A) > u) = \mathbf{P}(\{\mu_\infty^X(A) > u\} \circ \theta_{T_A}, T_A < \infty) = \sum_{a \in A} \mathbf{P}^a(\mu_\infty^X(A) > u) \mathbf{P}(T_A = T_a < \infty) \tag{2.30}$$

and since it is easy to see from its proof that (2.18) holds with \mathbf{P} replaced by \mathbf{P}^a for any $a \in A$, for some $c_1 = c_1(a)$ it follows that (2.18) also holds. This completes the proof of the lemma. \square

Remark. If A is replaced by $z + A$ for some fixed $z \in Z^d$, note from (1.2) that as matrices, $G_{z+A} = G_A$. Hence $\Lambda_{z+A} = \Lambda_A$.

3 Proof of Theorem 1.1

Given Lemma 2.2, Theorem 1.1 follows by the methods of [3, Section 7]. We spell out the details.

We first prove the lower bound for (1.4). To this end fix $a < \theta^{*-1} = 1/\log(\Lambda_A/(\Lambda_A - 1))$.

Let $k(n) = (\log n)^8$ and $N_n = \lfloor n/k(n) \rfloor$, and $t_{i,n} = ik(n)$ for $i = 0, \dots, N_n - 1$. Writing $X_s^t = X_{s+t} - X_t$ it follows that

$$\sup_{m \in [0, n]} \mu_n^X(X_m + A) \geq \max_{0 \leq i \leq N_n - 1} Z_i^{(n)},$$

where $Z_i^{(n)} = \mu_{k(n)}^{X_i^{t_i, n}}(A)$ are i.i.d. and by Lemma 2.2, for some $c > 0$ and all n large enough,

$$\mathbf{P}\left(\max_{0 \leq i \leq N_n - 1} Z_i^{(n)} \leq a \log n\right) \leq (1 - cn^{-a\theta^*})^{N_n} \leq e^{-cn^{-a\theta^*} N_n}.$$

Since $a\theta^* < 1$ this is summable, so that applying Borel-Cantelli, then taking $a \uparrow \theta^{*-1}$, we see that a.s.

$$\liminf_{n \rightarrow \infty} \sup_{m \in [0, n]} \frac{\mu_n^X(X_m + A)}{\log n} \geq \theta^{*-1}. \quad (3.1)$$

This gives the lower bound for (1.4).

For the upper bound, fix $a > \theta^{*-1}$. Note that for any $m \in [0, n]$

$$\begin{aligned} \mu_n^X(X_m + A) &= \sum_{j=0}^n \mathbf{1}_{X_m + A}(X_j) = \sum_{j=0}^n \mathbf{1}_A(X_j - X_m) \\ &= \sum_{j=0}^{m-1} \mathbf{1}_A(X_j - X_m) + \sum_{j=m}^n \mathbf{1}_A(X_j - X_m) \\ &\stackrel{\text{law}}{=} \sum_{j=1}^m \mathbf{1}_A(X'_j) + \sum_{j=0}^{n-m} \mathbf{1}_A(X''_j) \end{aligned} \quad (3.2)$$

where $\{X'_j, j = 0, 1, \dots\}$, $\{X''_j, j = 0, 1, \dots\}$ are two independent copies of $\{X_j, j = 0, 1, \dots\}$ and we have used the symmetry of X_1 . Using this and (2.28),

$$\begin{aligned} \mathbf{P}\left(\sup_{m \in [0, n]} \mu_n^X(X_m + A) \geq a \log n\right) &\leq \sum_{m=0}^n \mathbf{P}(\mu_n^X(X_m + A) \geq a \log n) \\ &\leq 2n\mathbf{P}(\mathcal{J} + \mathcal{J}' \geq a \log n) \leq c(\log n) n^{-(a\theta^* - 1)}. \end{aligned} \quad (3.3)$$

Thus letting $n_k = n^k$ for k sufficiently large that $k(a\theta^* - 1) > 2$, we see from applying Borel-Cantelli, then taking $a \downarrow \theta^{*-1}$, that a.s.

$$\limsup_{n \rightarrow \infty} \sup_{m \in [0, n^k]} \frac{\mu_{n^k}^X(X_m + A)}{\log n^k} \leq \theta^{*-1}.$$

The upper bound for (1.4) then follows by interpolation.

The lower bound for (1.3) follows immediately from (3.1). As for the upper bound in (1.3), we note that $\mu_n^X(x + A) = 0$ unless $X_m \in x + A$ for some $m \in [0, n]$. Thus the only relevant x 's in (1.3) are of the form $X_m - a$ for some $m \in [0, n]$ and $a \in A$. Thus

$$\sup_{x \in Z^d} \mu_n^X(x + A) = \sup_{m \in [0, n], a \in A} \mu_n^X(X_m - a + A). \quad (3.4)$$

Recalling Remark and the fact that A is a finite set, the upper bound for (1.3) now follows as in the proof of the upper bound for (1.4).

4 Examples

Proof of (1.5): When $A = \{0, y\}$ we have

$$G_A = \begin{pmatrix} G(0) & G(y) \\ G(y) & G(0) \end{pmatrix}.$$

The eigenvalues are $G(0) + G(y), G(0) - G(y)$ so that $\Lambda_A = G(0) + G(y) = G(0)(1 + t_y) = (1 + t_y)/\gamma_d$, where $t_y = \mathbf{P}(T_y < \infty)$, γ_d is the probability of no-return to the origin, and we have used the fact that $G(y) = t_y G(0)$. Then $1 - 1/\Lambda_A = 1 - \gamma_d/(1 + t_y)$.

We note that in the notation of Lemma 2.1, $h_1 = 1, h_2 = 0$ so that by (2.1)

$$\mathbf{P}(\mu_\infty^X(\{0, y\}) > u) = (1 - \gamma_d/(1 + t_y))^u \quad u = 1, 2, \dots \quad (4.1)$$

Proof of (1.6): We now consider the simple random walk, and for ease of notation consider first $d = 3$. Let $A = \{e_1, e_2, e_3, -e_1, -e_2, -e_3\} = S(0, 1)$, the (Euclidean) sphere in Z^3 of radius 1 centered at the origin. We have

$$G_{S(0,1)} = \begin{pmatrix} G(0) & G(e_1 - e_2) & G(e_1 - e_3) & G(2e_1) & G(e_1 + e_2) & G(e_1 + e_3) \\ G(e_2 - e_1) & G(0) & G(e_2 - e_3) & G(e_2 + e_1) & G(2e_2) & G(e_2 + e_3) \\ G(e_3 - e_1) & G(e_3 - e_2) & G(0) & G(e_3 + e_1) & G(e_3 + e_2) & G(2e_3) \\ G(2e_1) & G(e_1 + e_2) & G(e_1 + e_3) & G(0) & G(e_1 - e_2) & G(e_1 - e_3) \\ G(e_2 + e_1) & G(2e_2) & G(e_2 + e_3) & G(e_2 - e_1) & G(0) & G(e_2 - e_3) \\ G(e_3 + e_1) & G(e_3 + e_2) & G(2e_3) & G(e_3 - e_1) & G(e_3 - e_2) & G(0) \end{pmatrix}.$$

Using $G(x) = t_x G(0)$, where $t_x = \mathbf{P}(T_x < \infty)$, and symmetry which allows us to set $a =: t_{e_i \pm e_j}$ for $i \neq j$ and $b =: t_{2e_i}$ we can write

$$G_{S(0,1)} = G(0) \begin{pmatrix} 1 & a & a & b & a & a \\ a & 1 & a & a & b & a \\ a & a & 1 & a & a & b \\ b & a & a & 1 & a & a \\ a & b & a & a & 1 & a \\ a & a & b & a & a & 1 \end{pmatrix}.$$

It follows from the Perron-Frobenius Theorem that the largest eigenvalue is $\Lambda_{S(0,1)} = G(0)(1 + 4a + b)$ with eigenvector $(1, 1, 1, 1, 1, 1)$. Also, it is easy to see by symmetry that $\gamma_3 = \mathbf{P}(T_{e_i} = \infty)$. Then again by symmetry $\mathbf{P}(T_{e_1} = \infty) = \frac{4}{6}\mathbf{P}(T_{e_1-e_2} = \infty) + \frac{1}{6}\mathbf{P}(T_{2e_1} = \infty)$, i.e. $6\gamma_3 = 4\mathbf{P}(T_{e_1-e_2} = \infty) + \mathbf{P}(T_{2e_1} = \infty)$. Hence $\Lambda_{S(0,1)} = G(0)(1 + 4a + b) = G(0)6(1 - \gamma_3) = 6(1 - \gamma_3)/\gamma_3$. For the case of general $d \geq 3$, $G_{S(0,1)}$ is now a $2d \times 2d$ matrix, which is $G(0)$ times a matrix in which each row has a single entry of 1, a single entry of $b =: t_{2e_i}$ and $2d - 2$ entries of $a =: t_{e_i \pm e_j}$, where as before $t_x = \mathbf{P}(T_x < \infty)$. It is easy to see by symmetry that $\gamma_d = \mathbf{P}(T_{e_i} = \infty)$. Also, as before, it is easy to see by symmetry $\mathbf{P}(T_{e_1} = \infty) = \frac{(2d-2)}{2d}\mathbf{P}(T_{e_1-e_2} = \infty) + \frac{1}{2d}\mathbf{P}(T_{2e_1} = \infty)$, i.e. $2d\gamma_d = (2d - 2)\mathbf{P}(T_{e_1-e_2} = \infty) + \mathbf{P}(T_{2e_1} = \infty)$. Hence $\Lambda_{S(0,1)} = G(0)(1 + (2d - 2)a + b) = G(0)2d(1 - \gamma_d) = 2d(1 - \gamma_d)/\gamma_d$ for all $d \geq 3$.

We note that in the notation of Lemma 2.1, $h_1 = 1, h_j = 0, \forall j \neq 1$ so that by (2.1)

$$\mathbf{P}\left(\mu_\infty^X(S(0,1)) > u\right) = (1 - \gamma_d/2d(1 - \gamma_d))^u \quad u = 1, 2, \dots \quad (4.2)$$

Actually, (2.1) assumes that $0 \in A$ which doesn't hold here, but using (2.30) and symmetry we have that $\mathbf{P}\left(\mu_\infty^X(S(0,1)) > u\right) = \mathbf{P}^{e_1}\left(\mu_\infty^X(S(0,1)) > u\right)$ and (4.2) follows.

Proof of (1.7): We again consider the simple random walk. Let now $A = \{0\} \cup S(0,1) = B(0,1)$, the (Euclidean) ball in Z^3 of radius 1 centered at the origin. With $s = \mathbf{P}(T_{e_i} < \infty)$ and $\bar{s} = (s, \dots, s) \in R^{2d}$ we have

$$G_{B(0,1)} = G(0) \begin{pmatrix} 1 & \bar{s} \\ \bar{s}^t & M \end{pmatrix}$$

with M the $2d \times 2d$ matrix in the previous example. M is a self-adjoint matrix, and as mentioned the principal eigenvector is $\bar{1} = (1, \dots, 1) \in \mathbb{R}^{2d}$. If $u_i, i = 1, \dots, 2d - 1$ denote the other orthonormal eigenvectors of $G(0)M$ with eigenvalue $\lambda_i < \Lambda_{S(0,1)}$, then since they are orthogonal to $\bar{1}$ it is clear that $(0, u_i), i = 1, \dots, 2d - 1$ will give us $2d - 1$ orthonormal eigenvectors of $G_{B(0,1)}$ with eigenvalues $\lambda_i < \Lambda_{S(0,1)}$. The remaining (two) orthonormal eigenvectors must be of the form $(v, w\bar{1})$ and the corresponding eigenvalues will be $G(0)$ times those of the 2×2 matrix

$$L = \begin{pmatrix} 1 & 2ds \\ s & \Lambda \end{pmatrix}$$

where we abbreviate $\Lambda = \Lambda_{S(0,1)}/G(0) = 2d(1 - \gamma_d)$. The eigenvalues of L are

$$\frac{(1 + \Lambda) \pm \sqrt{(1 + \Lambda)^2 - 4(\Lambda - 2ds^2)}}{2} \quad (4.3)$$

so that

$$1/\Lambda_{B(0,1)} = \frac{2}{G(0)} \frac{(1 + \Lambda) - \sqrt{(1 + \Lambda)^2 - 4(\Lambda - 2ds^2)}}{4(\Lambda - 2ds^2)}. \quad (4.4)$$

Since $s = 1 - \gamma_d$ we have $\Lambda - 2ds^2 = 2d\gamma_d(1 - \gamma_d) = \gamma_d\Lambda$ we have

$$1/\Lambda_{B(0,1)} = \frac{(1 + \Lambda) - \sqrt{(1 + \Lambda)^2 - 4\gamma_d\Lambda}}{2\Lambda} \quad (4.5)$$

so that

$$\begin{aligned} 1 - 1/\Lambda_{B(0,1)} &= \frac{(\Lambda - 1) + \sqrt{(1 + \Lambda)^2 - 4\gamma_d\Lambda}}{2\Lambda} \\ &= \frac{(1 - 1/\Lambda) + \sqrt{(1 + 1/\Lambda)^2 - 4\gamma_d/\Lambda}}{2}. \end{aligned} \quad (4.6)$$

Setting $p = 1 - 1/\Lambda$ we can write this as

$$\begin{aligned} 1 - 1/\Lambda_{B(0,1)} &= \frac{p + \sqrt{(2 - p)^2 - 4\gamma_d/\Lambda}}{2} \\ &= \frac{p + \sqrt{p^2 + 4 - 4p - 4\gamma_d/\Lambda}}{2} \\ &= \frac{p + \sqrt{p^2 + 2/d}}{2}. \end{aligned} \quad (4.7)$$

We note that in the notation of Lemma 2.1, $h_j = 0$ for the $2d - 1$ orthonormal eigenvectors of the form $(0, u_i), i = 1, \dots, 2d - 1$ above. For the principle eigenvalue we have (4.7) and for the other ‘surviving’ eigenvalue the corresponding expression is $\frac{p - \sqrt{p^2 + 2/d}}{2}$. Hence by (2.1)

$$\mathbf{P} \left(\mu_\infty^X(B(0, 1)) > u \right) = h_1 \left(\frac{p + \sqrt{p^2 + 2/d}}{2} \right)^u + h_2 \left(\frac{p - \sqrt{p^2 + 2/d}}{2} \right)^u, \quad u = 1, 2, \dots, \quad (4.8)$$

where h_1, h_2 can be calculated in a straightforward manner. We observe that since $p < \sqrt{p^2 + 2/d}$, the expression in (4.8) is not a mixture of geometric random variables.

Now we prove Corollary 1.3. For any $y \in Z^d$ we have $t_y^2 < 1 - \gamma_d$, since t_y^2 is the probability that the random walk hits y and then returns to 0 in finite time which is obviously less than the probability $1 - \gamma_d$ that the random walk returns to zero in finite time. This implies $(1 + t_y - \gamma_d)^2 < (1 + t_y)^2(1 - \gamma)$ which in turn, implies

$$-1/\log(1 - \gamma_d/(1 + t_y)) < -2/\log(1 - \gamma_d),$$

and taking $\sup_{|y| \leq K}$ we obtain the Corollary 1.3.

5 The Brownian connection

Since R_K is a convolution operator on a bounded subset of \mathbb{R}^d with locally $L^1(\mathbb{R}^d, dx)$ kernel, it follows easily as in [4, Corollary 12.3] that R_K is a (symmetric) compact operator on $L^2(K, dx)$. Moreover, the Fourier transform relation $\int e^{i(x \cdot p)} u^0(x) dx = c|p|^{-2} > 0$ implies that R_K is strictly positive definite. By the standard theory for symmetric compact operators, R_K has discrete spectrum (except near 0) with all eigenvalues positive, and of finite multiplicity (see [7, Theorems VI.15, VI.16]). Moreover, if we use $(f, g)_{2,K}$ to denote the inner product in $L^2(K, dx)$, we have that $(f, R_K g)_{2,K} > 0$ for any non-negative, non-zero, f, g , so by the generalized Perron-Frobenius Theorem, see [8, Theorem XIII.43], the eigenspace corresponding to the largest eigenvalue, Λ_K^0 , is one dimensional.

Let $R_{K,\epsilon}$ be the operator on $L^2(K, dx)$ with kernel

$$u_{K,\epsilon}^0(x, y) = \sum_{z \neq z' \in \mathcal{L}_\epsilon(K)} u^0(z - z') 1_{e_\epsilon(z)}(x) 1_{e_\epsilon(z')}(y). \quad (5.1)$$

Since the sum is over disjoint sets, it can be checked easily that for any $1 < p < d/(d-2)$, $u_{K,\epsilon}^0(x, y)$ is bounded in $L^p(K \times K, dx dy)$ uniformly in $\epsilon > 0$:

$$\begin{aligned} \int_{K \times K} |u_{K,\epsilon}^0(x, y)|^p dx dy &= \sum_{z \neq z' \in \mathcal{L}_\epsilon(K)} \int_{e_\epsilon(z) \times e_\epsilon(z')} |u^0(z - z')|^p dx dy \\ &= c\epsilon^{2d} \sum_{z \neq z' \in \mathcal{L}_\epsilon(K)} \frac{1}{|z - z'|^{p(d-2)}} \\ &= c\epsilon^{2d-p(d-2)} \sum_{i \neq j \in \mathbb{Z}^d, |i|, |j| \leq k/\epsilon} \frac{1}{|i - j|^{p(d-2)}} \leq C. \end{aligned} \quad (5.2)$$

Also $u_{K,\epsilon}^0(x, y) \rightarrow u^0(x - y)$ as $\epsilon \rightarrow 0$ for all $x \neq y$. Hence, using (1.12)

$$\lim_{\epsilon \rightarrow 0} (f, R_{K,\epsilon} f)_{2,K} = (f, R_K f)_{2,K}, \quad \forall f \in C(K). \quad (5.3)$$

By [9],

$$G(x) = (1 + \delta(x))u^0(x), \quad \forall x \neq 0 \quad (5.4)$$

with $\delta(x)$ bounded and $\lim_{|x| \rightarrow \infty} \delta(x) = 0$ so that

$$\epsilon^{2-d} G(\epsilon^{-1}x) = (1 + \delta(\epsilon^{-1}x))u^0(x), \quad \forall x \in \mathcal{L}_\epsilon(K), x \neq 0. \quad (5.5)$$

Let $G_{K,\epsilon}$ be the operator on $L^2(K, dx)$ with kernel

$$v_{K,\epsilon}^0(x, y) = \sum_{z \neq z' \in \mathcal{L}_\epsilon(K)} \epsilon^{2-d} G(\epsilon^{-1}(z - z')) 1_{e_\epsilon(z)}(x) 1_{e_\epsilon(z')}(y). \quad (5.6)$$

Using (5.5), the same argument leading to (5.3) shows that

$$\lim_{\epsilon \rightarrow 0} (f, G_{K,\epsilon} f)_{2,K} = (f, R_K f)_{2,K}, \quad \forall f \in C(K). \quad (5.7)$$

Furthermore, since $G(0) < \infty$, if we let $\tilde{G}_{K,\epsilon}$ be the operator on $L^2(K, dx)$ with kernel

$$w_{K,\epsilon}^0(x, y) = \sum_{z, z' \in \mathcal{L}_\epsilon(K)} \epsilon^{2-d} G(\epsilon^{-1}(z - z')) 1_{e_\epsilon(z)}(x) 1_{e_\epsilon(z')}(y) \quad (5.8)$$

it follows from (5.7) that

$$\lim_{\epsilon \rightarrow 0} (f, \tilde{G}_{K,\epsilon} f)_{2,K} = (f, R_K f)_{2,K}, \quad \forall f \in C(K). \quad (5.9)$$

It now follows from [6, Theorem VIII.3.6,] that, if $\Lambda(\tilde{G}_{K,\epsilon})$ denotes the largest eigenvalue of the operator $\tilde{G}_{K,\epsilon}$

$$\lim_{\epsilon \rightarrow 0} \Lambda(\tilde{G}_{K,\epsilon}) = \Lambda(R_K) = \Lambda_K^0. \quad (5.10)$$

If f is any eigenvector for $\tilde{G}_{K,\epsilon}$ with eigenvalue $\zeta > 0$, it is clear that f is in the image of $\tilde{G}_{K,\epsilon}$ so that we can write

$$f(x) = \sum_{z \in \mathcal{L}_\epsilon(K)} 1_{e_\epsilon(z)}(x) f(z) \quad (5.11)$$

and the eigenvalue equation $\tilde{G}_{K,\epsilon} f = \zeta f$ becomes

$$1_{e_\epsilon(z)}(x) \sum_{z' \in \mathcal{L}_\epsilon(K)} \int_K w_{K,\epsilon}^0(x, y) 1_{e_\epsilon(z')}(y) f(z') dy = \zeta f(z), \quad \forall z \in \mathcal{L}_\epsilon(K). \quad (5.12)$$

Noting that the dy integration picks up a factor ϵ^d , this implies that

$$\sum_{z' \in \mathcal{L}_\epsilon(K)} \epsilon^2 G(\epsilon^{-1}(z - z')) f(z') = \zeta f(z), \quad \forall z \in \mathcal{L}_\epsilon(K). \quad (5.13)$$

Hence $\Lambda_{\epsilon^{-1}\mathcal{L}_\epsilon(K)} = \epsilon^{-2} \Lambda(\tilde{G}_{K,\epsilon})$. Together with (5.10) this completes the proof of Theorem 1.4.

References

1. Z. Ciesielski and S. J. Taylor, *First passage times and sojourn times for Brownian motion in space and the exact Hausdorff measure of the sample path*, Trans. Amer. Math. Soc. **103** (1962), 434–452.
2. A. Dembo, J. Rosen, Y. Peres and O. Zeitouni, *Thick points for spatial Brownian motion: multifractal analysis of occupation measure*, Ann. Probab., **28** (2000), 1-35.
3. P. Erdős and S. J. Taylor, *Some problems concerning the structure of random walk paths*, Acta Sci. Hung. **11** (1960), 137–162.
4. P. Halmos and V. Sunder, *Bounded Integral Operators on L^2 Spaces*, Springer-Verlag, New York, 1978.

5. G. Lawler, Notes on random walks, in preparation.
www.math.cornell.edu/~lawler/m778s04.html
6. T. Kato, *Perturbation Theory of Linear Operators*, second edition, Springer-Verlag, New York, 1976.
7. M. Reed and B. Simon, *Methods of Modern Mathematical Physics I: Functional Analysis*, Academic Press, New York, 1972.
8. M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators*, Academic Press, New York, 1978.
9. K. Uchiyama, *Green's functions for random walks on Z^N* , Proc. London Math. Soc. **77** (1998), 215–240.

Endre Csáki
Alfréd Rényi Institute of Mathematics
Hungarian Academy of Sciences
P.O. Box 127
H-1364 Budapest
Hungary
csaki@renyi.hu

Antónia Földes
Department of Mathematics
College of Staten Island, CUNY
2800 Victory Blvd., Staten Island
New York 10314
USA
afoldes@gc.cuny.edu

Pál Révész
Institut für Statistik und Wahrscheinlichkeitstheorie
Technische Universität Wien
Wiedner Hauptstrasse 8-10/107
A-1040 Vienna
Austria
revesz@ci.tuwien.ac.at

Jay Rosen
Department of Mathematics
College of Staten Island, CUNY
2800 Victory Blvd., Staten Island
New York 10314
USA
jrosen3@earthlink.net

Zhan Shi
Laboratoire de Probabilités UMR 7599
Université Paris VI
4 place Jussieu
F-75252 Paris Cedex 05
France
zhan@proba.jussieu.fr